

FOX RIVER

WATERSHED

INVENTORY AND ASSESSMENT

PREPARED BY

Robert A. Hrabik

Fisheries Management Biologist

October 1992

For additional information contact

Fisheries Regional Supervisor

Missouri Department of Conservation

2500 South Halliburton Kirksville, MO 63501

ACKNOWLEDGMENTS

This document was made possible through the diligent efforts of several individuals. Dwight Snead (SCS) provided invaluable land use information. Ken Brummett (MDC) provided support and insightful comments throughout the investigation. Randy Haydon and Kyle Reno (MDC) assisted in habitat and fish sampling. Randy Haydon also processed a significant amount of data on the computer. Mark Reigle (MDC) wrote the computer program MAPWORK, thereby saving a considerable amount of time and increasing our stream mapping accuracy. Dave Neuswanger (MDC) reviewed and revised the descriptive part of this document and prepared the planning section and executive summary. Rich Wehnes and Steve Eder (MDC) provided many helpful comments which were incorporated into the plan.

EXECUTIVE SUMMARY

The Fox River basin is a relatively small system of streams which drains 400 square miles in northeastern Missouri and southeastern Iowa upstream of the gage station on Fox River at Wayland, Missouri. Average annual discharge at the gage station is 258 cubic feet per second. The four largest streams in the basin are Fox River (52 miles long in Missouri), Little Fox River (24 miles in Missouri), Honey Creek (36 miles), and Sugar Creek (16 miles). Stream gradients average 3.0 feet per mile on Fox River, 5.0 feet per mile on Little Fox River, 6.7 feet per mile on Honey Creek, and 13.2 feet per mile on Sugar Creek.

Approximately 20% of basin stream mileage is channelized. While the Fox River itself is virtually unaltered, channelized reaches comprise 49% of Little Fox River, 41% of Sugar Creek, and 28% of Honey Creek.

Sedimentation is the only significant form of water pollution in the basin, but it threatens the integrity of the entire ecosystem. In 1984, the watershed was 63% cropland, 16% grassland, and 20% timber. The USDA Soil Conservation Service estimated that annual sediment delivery to the Fox and Wyaconda rivers averaged 3 tons per acre from the 483,780 acres which comprise the combined watersheds; this ranked ninth among 45 Missouri subbasins in rate of sediment delivery to stream channels. This sediment load equates to dumping 100,000 large truck loads of earth fill into these streams annually.

We have documented a reduction in Fox River base flow between the periods 1922-1952 and 1953-1980. Hydrological problems are most probably tied to land use practices which have diminished the moisture retention capacity of basin soils. These net adverse effects have been measurable despite a 5.8% increase in basin timber between 1939 and 1984, indicating that type of vegetative cover alone may not have as significant an effect on basin hydrology as the manner in which cover types are managed. From the 1950s through the 1980s, an increasing dependency on heavy machinery and chemical methods for producing crops has compacted the soil and reduced its organic matter content, thereby reducing its capacity to retain moisture.

The largely agricultural population of Clark County is generally unaware of the adverse effects that channelization, levee construction, riparian corridor clearing, and high-impact agriculture have had on basin streams.

In 1987, a Department of Conservation survey added 16 species to the annotated list of fishes known to the Missouri portion of the Fox River basin, which now number 52. Most fishes in our 1987 samples were widespread, tolerant species.

A statewide telephone survey revealed that 67% of Fox River anglers fished primarily for channel catfish. Our 1987 fish population surveys revealed that most channel catfish (84%) were small (<11 inches). Only 18% of 11-inch-and-larger channel catfish were of "quality" size (16 inches). We suspect that there is insufficient depth and current during much of the year to provide habitat suitable for quality-size channel catfish; they may migrate downstream to the Mississippi River prior to the onset of low-flow conditions. We also feel that migration of adult flathead catfish may significantly influence their density at any point in time. Our 1987 survey yielded only 28 flathead catfish, most small; yet several anglers have reported catching big flatheads during high-flow periods in late spring and early summer. Before we can manage catfish populations in the Fox River basin, we must know whether exploitable stocks are stable or transient. Also, we must learn which methods and times for sampling will provide meaningful information.

Relative to other stream basins in northeastern Missouri, Fox River receives very little attention by anglers or floaters. Boating and canoeing on all tributaries and most of Fox

River is hampered by shallow water, log jams, and low base flow. Even though recreational use of basin streams seems low relative to the availability of public stream frontage, there are some unique habitats which might be enjoyed if they were accessible.

Out 25-year strategic plan for the Fox River basin contains goals for the Fisheries Management Section of the Missouri Department of Conservation to improve aquatic habitat, maintain fish species richness and increase density of large sport fish, and increase appreciation for and accessibility to basin streams.

In order to improve aquatic habitat, we should do our best to prevent additional channelization projects, implement stream corridor management plans on public areas, convince basin farmers to engage in low-input sustainable agriculture and use acceptable methods for managing their riparian corridors, and cooperate with others in maintaining base flow at or above current levels.

In order to approach our goal for fish community integrity, we propose to maintain at least 50 native species of fish and to achieve "balanced" populations of channel and flathead catfish in basin streams. An important first step will be to learn enough about catfish migration patterns and catfish population survey methods to define seasonal parameters which may indicate whether "balance" exists in prairie streams.

In order to increase appreciation for and accessibility to basin streams, we should provide public access to the most unique and scenic reaches of basin streams and ensure that all potential stream anglers and floaters have access to information about recreational opportunities at these areas. To achieve this, we propose to amend the Department of Conservation's Stream Areas Acquisition Plan, develop a brochure, and generally facilitate public awareness and involvement with basin streams.

TABLE OF CONTENTS

<p style="text-align: center;"><i>WATERSHED LOCATION</i></p> <p style="text-align: center;"><i>LAND USE</i></p> <p>Historic and Recent Land Use</p> <p>Soil Conservation Projects</p> <p>Public Areas</p> <p>Corps of Engineers 404</p> <p>Jurisdiction</p> <p style="text-align: center;"><i>WATER QUALITY AND USE</i></p> <p>Beneficial Use Attainment</p> <p>Chemical Quality of Stream Flow</p> <p>Contamination Levels and Health Advisories</p> <p>Chronic Fish Kill Areas</p>	<p style="text-align: center;"><i>GEOLOGY / GEOMORPHOLOGY</i></p> <p>Physiographic Region</p> <p>Geology and Soils</p> <p>Watershed Area</p> <p>Channel Gradient</p> <p style="text-align: center;"><i>HYDROLOGY</i></p> <p>Precipitation</p> <p>USGS Gaging Stations</p> <p>Permanent / Intermittent Streams</p> <p>Average Annual Discharge</p> <p>7-day Q2 and Q10 Low Flows</p> <p>Dam and Hydropower Influences</p>
---	--

Water Use
Point Source Pollution
NonPoint Source Pollution

BIOTIC COMMUNITY

Fish Community Data
Aquatic Invertebrates
Threatened and Endangered
Species
Fish Stockings
Creel Survey Data

ANGLER GUIDE

GLOSSARY

LIST OF TABLES

HABITAT CONDITIONS

Channel Alterations
Unique Habitat
Improvement Projects
Stream Habitat Assessment

MANAGEMENT PROBLEMS/OPPORTUNITIES

LITERATURE CITED

RELATED INFORMATION

LIST OF FIGURES

List of Figures

Figure 1. Location of the Fox River watershed, Missouri and Iowa.

Figure nd. Location of the Fox River watershed within the natural divisions of Missouri and the landforms of Iowa.

Figure ge. Geology within the Fox River watershed, Missouri and Iowa.

Figure 4. Gradient plots of Fox River, Little Fox River, Honey Creek, and Sugar Creek.

Figure lu. Land Use/ Land Cover within the Fox River watershed, Missouri and Iowa (MORAP 1999, preliminary data).

Figure 6. Relationship of precipitation to streamflow in the Fox River.

Figure 7. Cumulative average flow of the Fox River, 1922-1988.

Figures 8a- 8c. Fox River duration curves.

Figure pa. Public areas in the Fox River Watershed in Missouri.

Figure gs. Gaging stations in the Fox River Watershed, in Missouri.

Figure ps. Point source pollution locations in the Fox River Watershed, in Missouri.

Figure fs. Fish sample locations in the Fox River Watershed, in Missouri.

Figure 13. Channel catfish length frequency in samples from the Fox River basin, 1987.

List of Tables

Table 1. Characteristics of stream channels in the Fox River Basin.

Table 2. Mean Gradient and Slope of Fox River and Its Major Tributaries.

Table 3. Changes in Forest Cover in the Fox River Basin, 1939-84.

Table 4. Public Stream Access and Frontage Areas in the Fox River Basin.

Table 5. Sport Fishery Characteristics of the Fox River and Other Northeastern Missouri Streams, Based Upon a Statewide Angler Telephone Survey, 1983-1986.

Table 6. Water Quality Parameters From Selected Sites in the Fox River Basin.

Table 7. Stream Habitat and Fish Sample Sites in the Fox River Basin, 1987.

Table 8. Habitat Parameters in the Fox River Basin, 1987.

Table 9. Annotated List and Status of Fishes Known to the Fox River Basin, Including Trophic, Geographical, and Ecological Affinities.

Table 10. Percentage Composition Within Sample Sites and Frequency of Occurrence Among All Sites of Fish Species Collected in the Fox River Basin, 1987.

Table 11. Age Structure of Channel and Flathead Catfish Collected from the Fox River Basin in 1987.

Table 12. Annotated List of Freshwater Mussels Sampled From the Fox River Basin in Missouri.